

DFT studies on hexaborate(2-) and heptaborate(2-) anions

Davies, Richard; Beckett, Michael; Jones, Charlotte

Phosphorus, Sulfur and Silicon and the Related Elements

DOI:

[10.1080/10426507.2015.1128923](https://doi.org/10.1080/10426507.2015.1128923)

Published: 01/04/2016

Peer reviewed version

[Cyswllt i'r cyhoeddiad / Link to publication](#)

Dyfyniad o'r fersiwn a gyhoeddwyd / Citation for published version (APA):

Davies, R., Beckett, M., & Jones, C. (2016). DFT studies on hexaborate(2-) and heptaborate(2-) anions. *Phosphorus, Sulfur and Silicon and the Related Elements*, 191(4), 633-637.
<https://doi.org/10.1080/10426507.2015.1128923>

Hawliau Cyffredinol / General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

Take down policy


If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

DFT studies on hexaborate(2-) and heptaborate(2-) anions

R. Andrew Davies,* Michael A. Beckett and Charlotte L. Jones

School of Chemistry, Bangor University, Bangor, Gwynedd, LL57 2UW, UK.

GRAPHICAL ABSTRACT



Abstract Bond counting rules allow chemists to unearth low-energy structures by sequentially sieving undesirable interactions whilst maximizing stabilizing interactions. The approach is applied to model hydrogen-bonding in polyborate anions.

Keywords Polyborate anions; hydrogen bonding; polyhydroxybenzoquinones

INTRODUCTION

Conformational analysis is a painstaking task even with advances in computing power. Bond counting rules (BCR) have been used to identify low-energy structures for B/C/N superhard materials,^{1,2} and doped fullerenes,³ based upon observations⁴ that stable structures had no B-B nor N-N bonds, whilst the number of C-C and B-N bonds were maximized. The BCR works well for B/C/N systems due to the large energetic differences (*ca.* 100-200 kJ mol⁻¹) between the weakest and strongest covalent bonds. It should be possible to extend BCRs to hydrogen-bonded systems,^{5,6} as typical hydrogen-bond energies (*ca.* 20 kJ mol⁻¹) are far greater than either thermal energy or computational error.

Received xx yyyy 2015; accepted xx yyyy 2015

We wish to thank Bangor University for funding.

Address correspondence to Dr. Andrew Davies, School of Chemistry, Bangor University, Bangor, Gwynedd, LL57 2UW, UK. Email: r.a.davies@bangor.ac.uk

Polyborate anions can form a myriad of structures including isolated rings or cages, 1-D chains, 2-D sheets and 3-D networks due to corner sharing linked triangular BO_3 and tetrahedral BO_4 units.⁷⁻¹¹

RESULTS AND DISCUSSION

Computational Methodology

Gas-phase B3LYP/6-311++G(*d,p*) DFT calculations were performed using *Gaussian09*.¹²

Hexaborate

The $[\text{B}_6\text{O}_7(\text{OH})_6]^{2-}$ anion consists of BO_3 triangles (Δ) & BO_4 tetrahedra (T) linked by vertices and a central 3-coordinate O atom to form three 6-membered rings containing 3- and 4-coordinate borons (Figure 1a) with a Christ and Clark description of $6:3\Delta + 3\text{T}$.⁷ The calculations suggest that *exo* OH groups attached to tetrahedral borons will preferentially orientate *anti*- to the ring B-O bonds. The most stable structures (Figure 2a) possess an unbroken, quasi-triangular cyclic network ($r_{\text{O-H}} = 0.967 \text{ \AA}$; $r_{\text{O...H}} = 2.25 \text{ \AA}$; $\angle_{\text{O-H...O}} = 135.7^\circ$; $r_{\text{O...O}} = 3.02 \text{ \AA}$) of cooperative intramolecular hydrogen bonds (IHBs). The cyclic network maximizes the number of stabilizing interactions in complete accord with bond counting rule approaches. Breaking an IHBs (so that two remain) *via* B-O bond rotation, raises the conformational energy by *ca.* 15 kJ mol^{-1} (Table 1), which is consistent with hydrogen bond energies.⁶ In complete contrast, rotation of the planar hydroxyl groups attached to a trigonal boron atom has a negligible energetic effect due to absence of H-bonding (no bond critical point) coupled with almost identical electrostatic attractions between the electropositive hydrogen of the OH group and the electronegative ring O atoms.

QTAIM analyses (Figure 2a) confirm the covalent nature of the OH groups with large ρ_{BCP} and -ve ∇^2 values (*ca.* 0.36 a.u. and -2.40 a.u. respectively). The closed-shell IHBs have much smaller ρ_{BCP} and ∇^2 values (*ca.* 0.02 a.u. and 0.05 a.u. respectively) reflecting their weaker nature. A 0.007 a.u. reduction in ρ_{BCP} , a 0.006 \AA elongation in r_{OH} and a *ca.* 100 cm^{-1} red shift in ν_{OH} are observed for *exo* vs planar OH groups, reflecting the loss in electron density that occurs during hydrogen bond formation.




Figure 1 Polyborate anion skeletal structures: (a) $[\text{B}_6\text{O}_7(\text{OH})_6]^{2-}$; (b) $[\text{B}_7\text{O}_9(\text{OH})_5]^{2-}$




Figure 2 QTAIM bond graph (AIM2000) and selected ρ_{BCP} values (a.u.): (a) $[\text{B}_6\text{O}_7(\text{OH})_6]^{2-}$; (b) $[\text{B}_7\text{O}_9(\text{OH})_5]^{2-}$.

Small red and yellow dots represent bond and ring critical points respectively.

Received xx yyyy 2015; accepted xx yyyy 2015

We wish to thank Bangor University for funding.

Address correspondence to Dr. Andrew Davies, School of Chemistry, Bangor University, Bangor, Gwynedd, LL57 2UW, UK. Email: r.a.davies@bangor.ac.uk

Table 1 Gas phase B3LYP/6-311++G(*d,p*) relative energies (kJ mol⁻¹) for the [B₆O₇(OH)₆]²⁻ hexaborate anion. Initial hydrogen atom orientations C = clockwise, A = anticlockwise and sites (a-f) relative to Figure 1a. Sixty four (2⁶) different combinations were generated giving rise to twelve unique enumerations.

Outer (Trigonal) OH groups			Inner (Tetrahedral) OH groups			ΔE (kJ mol ⁻¹)
a	b	c	d	e	f	
C	C	C	A	A	A	0
C	C	C	C	C	C	0
C	C	A	A	A	A	1
C	C	A	C	C	C	1
C	C	A	A	A	C	14
C	C	A	C	C	A	14
C	C	C	C	A	A	15
C	C	C	C	C	A	16
C	C	A	C	A	A	16
C	C	A	A	C	C	17
C	C	A	A	C	A	18
C	C	A	C	A	C	19

Table 2 Gas phase B3LYP/6-311++G(*d,p*) relative energies (kJ mol⁻¹) for the [B₇O₉(OH)₅]²⁻ heptaborate anion. Initial hydrogen atom orientations C = clockwise, A = anticlockwise and sites (a-e) relative to Figure 1b. Thirty two (2⁵) different combinations were generated giving rise to sixteen unique enumerations.

OH groups					ΔE (kJ mol ⁻¹)
a	b	c	d	e	
C	C	C	C	C	0
C	A	C	A	A	0
C	A	A	C	A	1
C	A	C	A	C	1
C	A	A	C	C	1
C	C	C	C	A	1
C	C	A	A	A	1
C	C	A	A	C	2
C	C	C	A	A	2
C	A	A	A	A	2
C	C	A	C	A	2
C	A	A	A	C	2
C	C	A	C	C	3
C	C	C	A	C	3
C	A	C	C	C	3
C	A	C	C	A	4

Received xx yyyy 2015; accepted xx yyyy 2015

We wish to thank Bangor University for funding.

Address correspondence to Dr. Andrew Davies, School of Chemistry, Bangor University, Bangor, Gwynedd, LL57 2UW, UK. Email: r.a.davies@bangor.ac.uk

Heptaborate

The $[\text{B}_7\text{O}_9(\text{OH})_5]^{2-}$ anion is related to $[\text{B}_6\text{O}_7(\text{OH})_6]^{2-}$ by condensing two *exo* OHs with a $\text{B}(\text{OH})_3$ unit, forming an additional 6-membered ring (Figure 1b). It has a Christ and Clark descriptor of $7:4\Delta + 3\text{T}$.⁷ An IHB is formed between the *exo*-OH and either O atom situated on the new ring (Figure 2b), giving rise to similar energies for all rotamers (Table 2). The weaker nature of the IHB ($r_{\text{O-H}} = 0.962 \text{ \AA}$; $r_{\text{O...H}} = 2.59 \text{ \AA}$; $\angle_{\text{O-H...O}} = 128.3^\circ$; $r_{\text{O...O}} = 3.27 \text{ \AA}$) is highlighted in the low ($< 0.01 \text{ a.u.}$) ρ_{BCP} value together with a much shorter BCP-RCP distance (almost to the point of BCP/NCP coalescence). A small (*ca.* 0.002 a.u.) reduction in ρ_{BCP} , together with a *ca.* 20 cm^{-1} red shift in ν_{OH} and negligible ($< 0.001 \text{ \AA}$) elongation in r_{OH} for *exo* vs planar hydroxyl groups confirms the weaker nature of the IHB. Rotation of the planar hydroxyl groups attached to trigonal boron atoms again have a negligible energetic effect due to absence of hydrogen bonding.

REFERENCES

1. Mattesini, M.; Matar, S. F. *Comput. Mater. Sci.*, **2001**, 20, 107-119.
2. Mattesini, M.; Matar, S.F. *Int. J. Inorg. Mater.*, **2001**, 3, 943-957.
3. Fan, X. F.; Zhu, Z.; Shen, Z. X.; Kuo, J.-L. *J. Phys. Chem. C*, **2008**, 112, 15691–15696.
4. Tateyama, Y.; Ogitsu, T.; Kusakabe, K.; Tsuneyuki, S.; Itoh, S. *Phys. Rev. B*, **1997**, 55, 10161
5. Beckett, M. A.; Davies, R. A.; Thomas, C.D. *Comput. Theor. Chem.*, **2014**, 1044, 74-79.
6. Beckett, M. A.; Coles, S. J.; Davies, R. A.; Horton, P. N.; Jones, C.L. *Dalton Trans.*, **2015**, 44, 7032-7040.
7. Christ, C. L.; Clark, J. R. *Phys. Chem. Miner.*, **1977**, 2, 59-87.
8. Farmer, J. B. *Adv. Inorg. Chem. Radiochem.*, **1982**, 25, 187-237.
9. Heller, G., *Top. Curr. Chem.*, **1986**, 131, 39-98.
10. Schubert, D. M.; *Struct. Bond.*, **2003**, 105, 1-40.
11. Belokoneva, E. L.; *Crystallogr. Rev.*, 2005, 11, 151-198.
12. Gaussian 09, Revision **D.01**, Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Scalmani, G.; Barone, V.; Mennucci, B.; Petersson, G. A.; Nakatsuji, H.; Caricato, M.; Li, X.; Hratchian, H. P.; Izmaylov, A. F.; Bloino, J.; Zheng, G.; Sonnenberg, J. L.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Vreven, T.; Montgomery, J. A., Jr.; Peralta, J. E.; Ogliaro, F.; Bearpark, M.; Heyd, J. J.; Brothers, E.; Kudin, K. N.; Staroverov, V. N.; Kobayashi, R.; Normand, J.; Raghavachari, K.; Rendell, A.; Burant, J. C.; Iyengar, S. S.; Tomasi, J.; Cossi, M.; Rega, N.; Millam, J. M.; Klene, M.; Knox, J. E.; Cross, J. B.; Bakken, V.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Martin, R. L.; Morokuma, K.; Zakrzewski, V. G.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Dapprich, S.; Daniels, A. D.; Farkas, Ö.; Foresman, J. B.; Ortiz, J. V.; Cioslowski, J.; Fox, D. J. Gaussian, Inc., Wallingford CT, 2009.